

Sub-Clover Selection Guide



SUB CLOVER VARIETY	RAINFALL (MM)																												
	300	325	350	375	400	425	450	475	500	525	550	575	600	625	650	675	700	725	750	775	800	825	850	875	900				
SUBTERRANEAN (SUITED TO BETTER DRAINED ACID SOILS)	NEW/ CURRENT	SF Tammin (M-88 SH-51%)																											
		SF Forbes (M-101 SH-33%)																											
		Bindoon																											
		SF Narrikup (M-126 SH-22%)																											
		SF Rosabrook (M-143 SH-29%)																											
	SUPERSEDED	Dalkeith																											
		Losa																											
		Seaton Park																											
		Campeda																											
		Mt Barker																											
YANNINICUMS (SUITED TO PERIODS OF WATER LOGGING)	NEW/ CURRENT	Monti																											
		SF Yanco (M-121 SH-45%)																											
		SF Rouse (M-131 SH-24%)																											
	SUPERSEDED	Trikkala/Riverina																											
		Gosse																											
BRACHYCALYCINUMS (SUITED TO ALKALINE OR CRACKING SOILS)	NEW/ CURRENT	Mawson																											
		SF Tarlee (M-130 SH-5%)																											
	SUPERSEDED	SF Antillo (M-136 SH-5%)																											
Mintaro										Clare																			
Antas																													

■ RAGT PRODUCTS
 ■ SUPERCEDED PRODUCTS
 ■ COMPETITOR PRODUCTS
 M FLOWERING MATURITY
 SH SEED HARDINESS

SELECTING THE BEST SUB-CLOVER VARIETY FOR YOUR SITUATION

The chart below can help you select varieties that have been bred for improvements in yield, pest and disease resistance, hardseededness and seed yield for regeneration. You should select the correct sub-species (subterranean, yanninicum, brachycalycinum or mix) for different soil types in paddock, then select the most appropriate variety for your rainfall. It is often advisable to mix 2-3 varieties to cover the fact that seasons and rainfall can vary by year. **For example:**

A mix of **Narrikup** and **Rosabrook** could be used in 600-650 mm rainfall to cover higher or lower rainfall years, or

A mix of **Rouse** and **Rosabrook** could be used where parts of a paddock can become waterlogged in some years but rainfall is reliable.

Contact Us



TERRITORY CONTACTS

Southern NSW

Daniel Zinga +61 437 323 630
danielzinga@ragt.au

North East VIC and Gippsland

Brett Tonkin +61 477 636 648
bretttonkin@ragt.au

Western VIC and SA

David Barnett +61 467 246 326
davidbarnett@ragt.au

Northern NSW

Matthew Duff +61 408 086 106
matthewduff@ragt.au

Northern VIC & TAS

David Gould +61 428 751 503
davidgould@ragt.au

Western Australia

Dale Thompson +61 427 903 943
dalethompson@ragt.au

National Sales Manager

Mark Palmer +61 499 701 715
markpalmer@ragt.au

Commercial Manager - Broadacre

David Peake +61 408 780 577
davidpeake@ragt.au

CUSTOMER SERVICE

Shepparton Office & Warehouse

13 Future Ct, Shepparton VIC 3630
info@ragt.au +61 3 5832 3800

Technical Product Development Manager - Forage

Richard Prusa +61 467 770 353
richardprusa@ragt.au

Technical Product Development Manager - Broadacre

David Leah +61 447 565 457
davidleah@ragt.au

 @RAGTAUSTRALIA

 @RAGTAU

 ragt.au