



Sub-Clover Selection Guide

SUB CLOVER VARIETY		RAINFALL (MM)																												
		300	325	350	375	400	425	450	475	500	525	550	575	600	625	650	675	700	725	750	775	800	825	850	875	900				
SUBTERRANEAN (SUITED TO BETTER DRAINED ACID SOILS)	NEW/ CURRENT	SF Tammin (<i>M</i> -88 <i>SH</i> -51%)																												
				SF Forbes (<i>M</i> -101 <i>SH</i> -33%)																										
							Bindoon																							
									SF Narrikup (<i>M</i> -126 <i>SH</i> -22%)																					
														SF Rosabrook (<i>M</i> -143 <i>SH</i> -29%)																
	SUPERSEDED		Dalkeith																											
				Losa																										
				Seaton Park																										
										Campeda																				
												Mt Barker																		
													Denmark/Goulburn																	
																		Leura												
NEW/ CURRENT				Monti																										
						SF Yanco (<i>M</i> -121 <i>SH</i> -45%)																								
											SF Rouse (<i>M</i> -131 <i>SH</i> -24%)																			
				Trikkala/Riverina																										
	SUPERSEDED									Gosse																				
													Larissa																	
																		Napier												
NEW/ CURRENT		Mawson																												
								SF Tarlee (<i>M</i> -130 <i>SH</i> -5%)																						
									SF Antillo (<i>M</i> -136 <i>SH</i> -5%)																					
					Mintaro																									
										Clare																				
SUPERSEDED									Antas																					

■ RAGT PRODUCTS
 ■ SUPERSEDED PRODUCTS
 ■ COMPETITOR PRODUCTS
 M FLOWERING MATURITY
 SH SEED HARDINESS

SELECTING THE BEST SUB-CLOVER VARIETY FOR YOUR SITUATION

The chart below can help you select varieties that have been bred for improvements in yield, pest and disease resistance, hardseededness and seed yield for regeneration. You should select the correct sub-species (subterranean, yanninicum, brachycalycinum or mix) for different soil types in paddock, then select the most appropriate variety for your rainfall. It is often advisable to mix 2-3 varieties to cover the fact that seasons and rainfall can vary by year. **For example:**

A mix of **Narrikup** and **Rosabrook** could be used in 600-650 mm rainfall to cover higher or lower rainfall years, or

A mix of **Rouse** and **Rosabrook** could be used where parts of a paddock can become waterlogged in some years but rainfall is reliable.

Contact Us



TERRITORY CONTACTS

Southern NSW

Clara Holmberg +61 437 323 630
claraholmberg@ragt.au

North East VIC and Gippsland

Brett Tonkin +61 477 636 648
bretttonkin@ragt.au

Western VIC and SA

Tahlia Ferguson +61 482 691 675
tahliaferguson@ragt.au

Northern NSW

Matthew Duff +61 408 086 106
matthewduff@ragt.au

Northern VIC & TAS

David Gould +61 428 751 503
davidgould@ragt.au

Western Australia

Dale Thompson +61 427 903 943
dalethompson@ragt.au

National Sales Manager

Mark Palmer +61 499 701 715
markpalmer@ragt.au

Commercial Manager - Broadacre

David Peake +61 408 780 577
davidpeake@ragt.au

Portfolio Lead

Richard Prusa +61 467 770 353
richardprusa@ragt.au

CUSTOMER SERVICE

Shepparton Office & Warehouse

13 Future Ct, Shepparton VIC 3630

info@ragt.au

+61 3 5832 3800



@RAGTAUSTRALIA



@RAGTAU



ragt.au

ragt.au



think
SOLUTIONS
think RAGT