


Sub-Clover Selection Guide



SUB CLOVER VARIETY	RAINFALL (MM)																											
	300	325	350	375	400	425	450	475	500	525	550	575	600	625	650	675	700	725	750	775	800	825	850	875	900			
SUBTERRANEAN (SUITED TO BETTER DRAINED ACID SOILS)	NEW/ CURRENT	SF Tammin (M-88 SH-51%)																										
		SF Forbes (M-101 SH-33%)																										
		Bindoon																										
		SF Narrikup (M-126 SH-22%)																										
		SF Rosabrook (M-143 SH-29%)																										
	SUPERSEDED	Dalkeith																										
		Losa																										
		Seaton Park																										
		Campeda																										
		Mt Barker																										
YANNINICUMS (SUITED TO PERIODS OF WATER LOGGING)	NEW/ CURRENT	Monti																										
		SF Yanco (M-121 SH-45%)																										
		SF Rouse (M-131 SH-24%)																										
	SUPERSEDED	Trikkala/Riverina																										
		Gosse																										
BRACHYCALYCINUMS (SUITED TO ALKALINE OR CRACKING SOILS)	NEW/ CURRENT	Mawson																										
		SF Tarlee (M-130 SH-5%)																										
	SUPERSEDED	SF Antillo (M-136 SH-5%)																										
Mintaro																												
Clare																												
Antas																												

■ RAGT PRODUCTS
 ■ SUPERSEDED PRODUCTS
 ■ COMPETITOR PRODUCTS
 M FLOWERING MATURITY
 SH SEED HARDINESS

SELECTING THE BEST SUB-CLOVER VARIETY FOR YOUR SITUATION

The chart below can help you select varieties that have been bred for improvements in yield, pest and disease resistance, hardseededness and seed yield for regeneration. You should select the correct sub-species (subterranean, yanninicum, brachycalycinum or mix) for different soil types in paddock, then select the most appropriate variety for your rainfall. It is often advisable to mix 2-3 varieties to cover the fact that seasons and rainfall can vary by year. **For example:**

A mix of **Narrikup** and **Rosabrook** could be used in 600-650 mm rainfall to cover higher or lower rainfall years, or

A mix of **Rouse** and **Rosabrook** could be used where parts of a paddock can become waterlogged in some years but rainfall is reliable.

Contact Us



TERRITORY CONTACTS

Northern Victoria

Ian Carkeek +61 474 103 946
iancarkeek@ragt.au

Gippsland & Tasmania

Linda Ferguson +61 477 636 648
lindaferguson@ragt.au

Southern New South Wales

Georgie Constance +61 437 323 630
georgieconstance@ragt.au

Northern NSW

Matthew Duff +61 408 086 106
matthewduff@ragt.au

South Australia & Western Victoria

Mark Palmer +61 482 691 675
markpalmer@ragt.au

Western Australia

Dale Thompson +61 427 903 943
dalethompson@ragt.au

National Sales Manager

Mark Palmer
+61 499 701 715
markpalmer@ragt.au

Broadacre Business Development Manager

Kirk Brehaut
+61 459 858 845
kirkbrehaut@ragt.au

Broadacre Business Development Manager WA

David Peake
+61 408 780 577
davidpeake@ragt.au

Portfolio Lead

Richard Prusa
+61 467 770 353
richardprusa@ragt.au

CUSTOMER SERVICE

Shepparton Office & Warehouse

13 Future Ct, Shepparton VIC 3630

info@ragt.au

+ 61 3 5832 3800

 @RAGTAUSTRALIA

 @RAGTAU

 ragt.au